arXiv:1302.4593v2 [math.DG] 27 Feb 2013

SINGULAR RIEMANNIAN FOLIATIONS AND APPLICATIONS TO POSITIVE AND NONNEGATIVE CURVATURE

FERNANDO GALAZ-GARCIA* AND MARCO RADESCHI

Abstract. We study singular Riemannian foliations whose leaves are homeomorphic to Bieberbach manifolds. These foliations, which we call *B-foliations*, generalize isometric torus actions on Riemannian manifolds. We apply our results to the classification problem of closed, simply connected Riemannian 4- and 5-manifolds with positive or nonnegative curvature.

Contents

1. Introduction 1
2. Preliminaries 4
3. B-foliations 7
4. Euler characteristic of B-foliations 13
5. B-foliations of codimension 1 on simply connected manifolds 15
6. B-foliations of codimension 2 on simply connected manifolds 17
7. Curvature and B-foliations of codimension 2 20
8. Curvature and singular Riemannian foliations by circles 27
References 31

1. Introduction

The study of effective smooth torus actions on compact, smooth manifolds has a rich and long tradition in the the theory of smooth transformation groups (cf. [5, 11, 12, 28, 30, 34, 35]). In Riemannian geometry, starting with Hsiang and Kleiner's topological classification of closed Riemannian 4-manifolds of positive (sectional) curvature with an effective

---

Date: February 28, 2013.

2000 *Mathematics Subject Classification.* 53C20, 53C23, 53C12, 57R30.

Key words and phrases. codimension 2, torus, singular Riemannian foliation, positive curvature, nonnegative curvature, 4-manifold, 5-manifold.

*The author is part of SFB 878: *Groups, Geometry & Actions*, at the University of Münster.

isometric circle action [26], isometric actions of tori have been successfully used to obtain classification results on compact Riemannian manifolds with positive or nonnegative curvature and large isometry groups (cf. [16, 17, 21, 29, 26, 42, 44, 46, 48]).

The present paper's main contribution is the observation that several results on smooth torus actions, and on isometric torus actions on Riemannian manifolds with positive or nonnegative curvature, hold under less restrictive conditions which do not involve the existence of a torus action. Indeed, many of these results do not hold because of the presence of a torus action, but rather because the orbit decomposition of the manifold has the structure of a *singular Riemannian foliation* whose leaves are diffeomorphic to flat tori of possibly different dimensions. To make this statement precise, we introduce a special class of singular Riemannian foliations, *B-foliations*, which generalize isometric torus actions on Riemannian manifolds. Roughly speaking, a B-foliation $(M, \mathcal{F})$ is a partition of a Riemannian manifold M into connected submanifolds, called the *leaves* of $\mathcal{F}$, all of which are homeomorphic to some flat manifold and are at a constant distance from each other.

The fact that B-foliations are more general than isometric torus actions is clear for several reasons. On the one hand, the leaves need not be tori. On the other hand, even when the leaves are diffeomorphic to tori, there might not be a global torus action on the manifold; this occurs, for example, when the distribution of the tangent spaces of the torus leaves is not orientable.

We point out that another generalization of isometric torus actions are the so-called *F-structures*, introduced by Cheeger and Gromov [7, 8]. These structures are, roughly speaking, generalized local torus actions and play a central role in the Cheeger-Fukaya-Gromov theory of collapsed Riemannian manifolds with bounded curvature (cf. [6, 14]). Although one may expect F-structures and B-foliations to be related, it is not clear to what extent the two concepts overlap, since neither one implies the other. For example, B-foliations with exotic torus leaves cannot be generated by F-structures, and certain F-structures among those that are not pure (see [45] for definitions) do not generate a B-foliation.

In this paper we focus our attention on B-foliations on simply connected manifolds and extend to B-foliations several basic results on torus actions on smooth compact manifolds. In particular, we prove:

Theorem A. *The regular leaves of a B-foliation on a simply connected manifold are homeomorphic to tori.*

Theorem B. *Let $(M, \mathcal{F})$ be a B-foliation and let $\Sigma_0 \subseteq M$ denote the stratum of 0-dimensional leaves. Then $\chi(\Sigma_0) = \chi(M)$.*

Theorem C. *Let $(M^{n+2}, \mathcal{F}^n)$ be a codimension 2 B-foliation on a compact, simply connected manifold M^{n+2} with $n \geq 1$. Then either $M = \mathbb{S}^3$ and $\mathcal{F}$ is given by a weighted Hopf action, or the following hold:*

(1) *The leaf space* $B = M/\mathcal{F}$ *is homeomorphic to a* 2*-disk, the interior of* B *is smooth, and the boundary* ∂B *consists of at least* n *totally geodesic segments meeting at an angle of* $\pi/2$.

(2) *Let* L_0 *be a generic leaf and* L_1 *a singular leaf. Then there is a submersion* $L_0 \to L_1$*, with fiber* $\mathbb{S}^1$ *if* L_1 *belongs to a geodesic in* ∂B*, or with fiber* T^2*, if* L_1 *belongs to a vertex of* ∂B.

It is worth pointing out that we assume the leaves of a B-foliation to be homeomorphic to flat manifolds, but not necessarily diffeomorphic. Therefore, Theorems A, B and C also hold for B-foliations where the leaves carry exotic smooth structures. Nevertheless, we do not know of any non-trivial examples of such foliations. It would be interesting to find B-foliations by exotic tori on simply connected manifolds and, in particular, on spheres.

Recall that a Riemannian manifold (M, g) is said to have *quasi-positive curvature* if (M, g) has nonnegative (sectional) curvature and a point with strictly positive curvature. As an application of our results, we classify compact, simply connected 4- and 5-manifolds with quasi-positve or nonnegative curvature and a codimension 2 B-foliation, and compact, simply connected 4-manifolds of positive or nonnegative curvature and a codimension 3 foliation by circles:

Theorem D. *Let* (M^n, g) *be a compact, simply connected Riemannian* n*-manifold with quasi-positive curvature supporting a codimension* 2 *B-foliation.*

(1) *If* $n = 4$*, then* M^4 *is diffeomorphic to* $\mathbb{S}^4$ *or* $\mathbb{C}P^2$.

(2) *If* $n = 5$*, then* M^5 *is diffeomorphic to* $\mathbb{S}^5$.

Theorem E. *Let* (M^n, g) *be a compact, simply connected Riemannian* n*-manifold with nonnegative curvature and a codimension* 2 *B-foliation.*

(1) *If* $n = 4$*, then* M^4 *is diffeomorphic to* $\mathbb{S}^4$, $\mathbb{C}P^2$ *or homeomorphic to* $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ *or* $\mathbb{S}^2 \times \mathbb{S}^2$.

(2) *If* $n = 5$*, then and* M^5 *is diffeomorphic to* $\mathbb{S}^5$ *or to one of the two* $\mathbb{S}^3$*-bundles over* $\mathbb{S}^2$.

Theorem F. *Let* M *be a compact, simply connected Riemannian* 4*-manifold with a singular Riemannian foliation by circles. Then the foliation is induced by a smooth circle action and the following hold:*

(1) *If* M *has positive curvature, then* M *is diffeomorphic to* $\mathbb{S}^4$ *or* $\mathbb{C}P^2$.

(2) *If M has nonnegative curvature, then M is diffeomorphic to* $\mathbb{S}^4$, $\mathbb{C}P^2$ *or* $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ *or* $\mathbb{S}^2 \times \mathbb{S}^2$.

Theorems D, E and F extend results in [26, 29, 46, 17] to the case of B-foliations. In all cases, the classification is up to diffeomorphism, except for (1) in Theorem E. This is because in this case we do not know whether the B-foliation is induced by a smooth torus action.

Our paper is structured as follows. In Section 2 we recall some basic facts on singular Riemannian foliations. In Section 3 we introduce the concept of B-foliation, we show that the infinitesimal foliation at any point of a manifold with a B-foliation is also a B-foliaton (see Subsection 2.1.4) and prove Theorem A. In Section 4 we prove Theorem B. The main ingredient in the proof is the existence of a good open cover on the quotient space of a closed singular Riemannian foliation. In Section 5 we classify codimension 1 B-foliations on simply connected manifolds. In Section 6 we prove Theorem C. In Section 7 we use Theorems A and C, together with the Mayer-Vietoris spectral sequence, to prove Theorems D and E. Finally, in Section 8 we prove Theorem F. Secondary results, possibly of independent interest, are distributed throughout the paper.

Acknowledgements. The second named auhor thanks Xiaoyang Chen for some initial conversations. Both authors wish to acknowledge the hospitality of the Mathematisches Forschungsinstitut Oberwolfach, where the work presented on this paper was initiated. The authors also wish to thank Wolfgang Ziller for helpful comments on a first version of this paper and Alexander Lytchak for helpful suggestions which led to the proofs of Theorems A and 3.6.

2. Preliminaries

In this section we collect some background material on singular Riemannian foliations.

2.1. **Riemannian foliations.**

In this subsection we recall some basic facts about singular Riemannian foliations. We refer the reader to [32, 1] for further results on the theory.

2.1.1. *Singular Riemannian foliations.* A *transnormal system* $\mathcal{F}$ on a Riemannian manifold M is a decomposition of M into smooth, complete, injectively immersed connected submanifolds, called *leaves*, such that every geodesic emanating perpendicularly to one leaf remains perpendicular to all leaves. A transnormal system $\mathcal{F}$ is called a *singular Riemannian foliation* if there are smooth vector fields X_i on M such that, for each point $p \in M$, the tangent space T_pL_p to the leaf L_p through p is given as the span of the vectors $X_i(p) \in T_pM$. We will call the quotient space $M/\mathcal{F}$ the *leaf space*, and will occasionally denote it by M^*. We will let $\pi : M \to M/\mathcal{F}$ be the leaf projection map. The pair $(M, \mathcal{F})$ will denote a singular Riemannian foliation $\mathcal{F}$ on a Riemannian manifold M. Slightly abusing notation, we will also refer to the pair $(M, \mathcal{F})$ as a singular Riemannian foliation.

A singular Riemannian foliation $\mathcal{F}$ will be called *closed* if all its leaves are closed in M; the foliation will be called *locally closed* at $x \in M$ if, for some neighborhood U of x, the restriction of $\mathcal{F}$ to U is closed, i.e. connected components of the intersection of the leaves of $\mathcal{F}$ with U are closed in U. If $\mathcal{F}$ is locally closed at x, then the local quotient $U/\mathcal{F}$ is a well defined Alexandrov space of curvature locally bounded from below. Similarly, if $\mathcal{F}$ is closed and M is complete, then the quotient space $M/\mathcal{F}$ is an Alexandrov space of curvature locally bounded below.

2.1.2. *Group actions.* As group actions will appear throughout our work, let us fix some notation before proceeding. Given a Lie group G acting (on the left) on a smooth manifold M, we denote by $G_p = \{\, g \in G : gp = p \,\}$ the *isotropy group* at $p \in M$ and by $Gp = \{\, gp : g \in G \,\} \simeq G/G_p$ the *orbit* of p. The *ineffective kernel* of the action is the subgroup $K = \cap_{p \in M} G_p$. We say that G acts *effectively* on M if K is trivial. The action is *free* if every isotropy group is trivial. Given a subset $A \subset M$, we will denote its image in M/G under the orbit projection map by A^*. When convenient, we will also denote the orbit space M/G by M^*.

Example 2.1 (Isometric Lie group actions)**.** Perhaps the most familiar example of a singular Riemannian foliation is the foliation induced by an (effective) isometric action of a Lie group G on a Riemannian manifold M. In this case, the foliation is given by the orbits of the action, and we say that the foliation is a *homogeneous foliation.* If G is compact, then the foliation is closed, and it is locally closed if and only if all the slice representations $G_p \to O(\nu_p(Gp))$ have compact image.

Remark 2.2. We will sometimes denote a homogeneous foliation, given by the action of a Lie group G, by (M, G), provided the G-action is understood.

2.1.3. *Stratification.* Let M be a Riemannian manifold with a singular Riemannian foliation $\mathcal{F}$. The *dimension* of $\mathcal{F}$, denoted by $\dim \mathcal{F}$, is the maximal dimension of its leaves. The *codimension* of $\mathcal{F}$ is, by definition,

$$\operatorname{codim}(\mathcal{F}, M) = \dim M - \dim \mathcal{F}.$$

For $k \leq \dim \mathcal{F}$, define

$$\Sigma_{(k)} = \{\, p \in M : \dim L_p = k \,\}.$$

Every connected component C of the set $\Sigma_{(k)}$ is an embedded (possibly non complete) submanifold of M and the restriction of $\mathcal{F}$ to C is a Riemannian foliation. Given $p \in M$, let Σ^p be the connected component of $\Sigma_{(k)}$ through p, where $k = \dim L_p$. We will refer to the decomposition of M into the submanifolds Σ^p as the *canonical stratification* of M.

The subset $\Sigma_{(\dim \mathcal{F})}$ is open, dense and connected in M; it is called the *regular stratum* of M. It will be denoted by M_0 and its points will be called *regular points.* If $M_0 = M$ we say that the foliation is *regular.* All other strata Σ^p have codimension at least 2 in M and are called *singular strata.* For any singular stratum Σ^p, we have

$$\operatorname{codim}(\mathcal{F}, \Sigma^p) < \operatorname{codim}(\mathcal{F}, M).$$

2.1.4. *Infinitesimal singular Riemannian foliations.* Let M be a Riemannian manifold with a singular Riemannian foliation $\mathcal{F}$. Given a point $p \in M$ and some small $\epsilon > 0$, let $S_p = \exp_p(\nu_p L_p) \cap B_\epsilon(p)$ be a *slice* through p, where $B_\epsilon(p)$ is the distance ball of radius ϵ around p. The foliation $\mathcal{F}$ induces a foliation $\mathcal{F}|_{S_p}$ on S_p by letting the leaves of $\mathcal{F}|_{S_p}$ be the connected components of the intersection between S_p and the leaves of $\mathcal{F}$. The foliation $(S_p, \mathcal{F}|_{S_p})$ may not be a singular Riemannian foliation with respect to the induced metric on S_p. Nevertheless, the *pull-back* foliation $\exp_p^*(\mathcal{F})$ is a singular Riemannian foliation on $\nu_p L_p \cap B_\epsilon(0)$ equipped with the Euclidean metric (cf. [32, Proposition 6.5]), and it is invariant under homotheties fixing the origin (cf. [32, Lemma 6.2]). In particular, it is possible to extend $\exp^*(\mathcal{F})$ to all of $\nu_p L_p$, giving rise to a singular Riemannian foliation $(\nu_p L_p, \mathcal{F}_p)$ called the *infinitesimal foliation* of $\mathcal{F}$ at p.

Notice that $0 \in \nu_p(L_p)$ is always a leaf of the infinitesimal foliation $\mathcal{F}_p$. By definition, leaves stay at a constant distance from each other, in particular every leaf stays in some distance sphere around the origin, and it makes sense to consider the infinitesimal foliation restricted to the unit sphere. Since the infinitesimal foliation is invariant under homothetic transformations, it can be reconstructed from its own restriction to the unit sphere. Taking this into account, we will sometimes refer to $(\nu_p^1 L_p, \mathcal{F}_p)$ also as the infinitesimal foliation at p and shall write $(\mathbb{S}_p^\perp, \mathcal{F}_p)$.

Given two points p_1, p_2 in some leaf L, the corresponding infinitesimal foliations $(\mathbb{S}_{p_1}^\perp, \mathcal{F}_{p_1})$, $(\mathbb{S}_{p_2}^\perp, \mathcal{F}_{p_2})$ are foliated isometric, in the sense that there is a (non-canonical) linear isometry $\mathbb{S}_{p_1}^\perp \to \mathbb{S}_{p_2}^\perp$ preserving the foliation. Moreover, these foliations can be glued together to give a foliation on $\nu^1(L)$ in the following sense: If one identifies $\nu^1(L)$ via the normal exponential map with $\partial \mathrm{Tub}_\varepsilon(L)$, the boundary of an ε-tubular neighborhood of L, then the intersections of leaves in $\mathcal{F}$ with with $\mathbb{S}_p^\perp$ are exactly the leaves in $\mathcal{F}_p$. In particular, if L' is a leaf in $\partial \mathrm{Tub}_\varepsilon(L) \simeq \nu^1 L$, then L' is a union of infinitesimal leaves. Moreover, if $p \in L$ and $q \in L'$ can be written as $q = \exp_p \varepsilon v$, $v \in \mathbb{S}_p^\perp$, then the connected components of a fiber of p under the metric projection $L' \to L$ (which is a submersion, cf. [32, Lemma 6.1]) are given by $\mathcal{L}_v$, where $\mathcal{L}_v \in \mathcal{F}_p$ are diffeomorphic to the infinitesimal leaf of $\mathbb{S}_p^\perp$ passing through v. Therefore, there is a fibration

$$\mathcal{L}_v \to L_q \to \bar{L}_p \tag{2.1}$$

for some finite cover $\bar{L}_p \to L_p$.

Remark 2.3. Since we may consider the leaf space M^* as an Alexandrov space (with curvature bounded below), let us quickly recall the procedure to compute the space of directions Σ_{p^*} at a point $p^* \in M^*$. Let $p \in M$ be a point in the preimage of p^* and let $\mathbb{S}_{p^*} = \mathbb{S}_p^{\perp}/\mathcal{F}_p$ be the quotient of the infinitesimal foliation $(\mathbb{S}_p^{\perp}, \mathcal{F}_p)$. The fundamental group $\pi_1(L_p)$ acts on $\mathbb{S}_{p^*}$ by isometries via the so-called *holonomy action* and Σ_{p^*} is isometric to $\mathbb{S}_{p^*}/\pi_1(L_p)$. Given $v \in \mathbb{S}_p^{\perp}$ with image $v^* \in \mathbb{S}_{p^*}$, let H be the subgroup of $\pi_1(L_p)$ fixing v^*. Then, in fibration (2.1), the cover $\bar{L}_p$ is $\tilde{L}_p/H$, where $\tilde{L}_p$ is the universal cover of L_p.

Example 2.4. Let (M, G) be a homogeneous foliation. Given a point $p \in M$, the connected component G_p^0 of the isotropy group G_p acts on $\nu_p(Gp)$ by isometries, via the so-called *slice representation*. In this case, the infinitesimal foliation $\mathcal{F}_p$ is the homogeneous foliation given by the orbits of G_p^0 on $\nu_p(Gp)$. Given $q \in M$ close to p, with isotropy $G_q < G_p$, the projection (2.1) is the projection

$$G_p^0/G_q \to G/G_q \to G/G_p^0,$$

where G/G_p^0 is a cover of the orbit G/G_p though p.

3. B-FOLIATIONS

We now introduce *B-foliations*, which are the main object of study in our paper. Recall that a *Biebarbach manifold* is a manifold diffeomorphic to $\mathbb{R}^n/G$, where G is a discrete group of Euclidean isometries acting freely and cocompactly on $\mathbb{R}^n$. These groups are called *Bieberbach groups*. Abstractly, Bieberbach groups can be characterized as torsion-free groups with a normal finite index abelian subgroup (cf. [50]). In particular, every subgroup of a Bieberbach group is a Bieberbach group. Every Bieberbach manifold is compact, has no boundary and admits a flat Riemannian metric.

Remark 3.1. We shall use the fact that any acyclic manifold N with fundamental group isomorphic to a Bieberbach group G must be homeomorphic to a Bieberbach manifold. To see this, observe first that N must be homotopy equivalent to $\mathbb{R}^n/G$, since both are models for $K(G, 1)$. It then follows from the solution of the Borel conjecture for flat manifolds (cf. [10, Section 4] and [4]) that N and $\mathbb{R}/G$ must be homeomorphic. The manifolds N and $\mathbb{R}^n/G$ may not be diffeomorphic, as illustrated by the examples of 5-tori which are mutually homeomorphic but not diffeomorphic (cf. [25]).

Definition 3.2. A closed singular Riemannian foliation $(M, \mathcal{F})$ is a *B-foliation* if every leaf is homeomorphic to some Bieberbach manifold.

Remark 3.3. In the preceding definition we do not assume that the leaves are flat with the induced Riemannian metric.

Example 3.4. Every isometric torus action on a Riemannian manifold induces a (homogeneous) B-foliation. Bundles whose fibers are homeomorphic to Bieberbach manifolds are also examples of B-foliations.

Example 3.5 (Non-homogeneous B-foliations)**.** The simplest way to construct non-homogeneous B-foliations with regular leaves homeomorphic to tori is to take the Riemannian product of a Riemannian manifold and an exotic torus. As the leaves are exotic tori, they cannot correspond to the orbits of an isometric torus action. One can also construct non-homogeneous B-foliations whose leaves are diffeomorphic to flat tori in the following way. Let B be a compact smooth manifold with infinite cyclic fundamental group, for example, $B = \mathbb{S}^1$, and let T^n be a standard n-dimensional torus. By a result of Hatcher [24], for n large enough, the diffeomorphism group $\mathrm{Diff}(T^n)$ has infinitely many connected components. Consider a representation

$$\pi_1(B) \to \mathrm{Diff}(T^n)$$

which maps a generator g of $\pi_1(B)$ to a connected component of $\mathrm{Diff}(T^n)$ which is not a "standard" one, i.e. one not containing any linear diffeomorphism of T^n. Let $\tilde{B}$ be the universal cover of B and let $\pi_1(B)$ act diagonally on the product $\tilde{B} \times T^n$. This action is free and, taking the quotient, we obtain a fiber bundle

$$T^n \to (\tilde{B} \times T^n)/\pi_1(B) \to B.$$

The total space of the bundle is B-foliated. On the other hand, since the structure group is not linear, the B-foliation cannot be homogeneous .

3.1. **The infinitesimal foliation of a B-foliation.**

In the case of B-foliations, the total space and the base in fibration (2.1) are homeomorphic to Bieberbach manifolds. The following general result shows that for B-foliations such fibrations are somewhat rigid.

Theorem 3.6. *Let $F \to M \to N$ be a fibration. If M is homeomorphic to a Bieberbach manifold, then F and N are both homeomorphic to Bieberbach manifolds.*

Proof. By Remark 3.1 it suffices to show that F and N are acyclic and $\pi_1(F)$ and $\pi_1(N)$ are Bieberbach groups.

Consider the fibration between the universal covers $F \to \tilde{M} \to \tilde{N}$. Since $\tilde{M}$ is contractible and $\tilde{N}$ is simply connected, we can apply the Serre spectral sequence with integral coefficients, and from it we obtain that F and N are contractible. In fact, if $H^*(F)$ has cohomological dimension a, and $H^*(\tilde{N})$ has cohomological dimension b, then $H^*(\tilde{M})$ would have cohomological dimension $a+b$ and this has to be 0. Therefore, N is acyclic, and from the exact sequence in homotopy, so is F.

From the same exact sequence, we have

$$1 \to \pi_1(F) \to \pi_1(M) \to \pi_1(N) \to 1,$$

where $\pi_1(M)$ is a Bieberbach group, i.e. a torsion free group with a finite index normal abelian subgroup. Since $\pi_1(F)$ is a subgroup of a Bieberbach group, it is again a Bieberbach group and therefore F is homeomorphic to a flat manifold.

We now prove that $\pi_1(N)$ is a Bieberbach group. First, we show that $\pi_1(N)$ is torsion free. Suppose this is not the case. Then there is a finite cyclic subgroup $\mathbb{Z}_k$ acting freely on the contractible manifold $\tilde{N}$. It follows that $\tilde{N}/\mathbb{Z}_k$ is a $K(\mathbb{Z}_k, 1)$, which contradicts the fact that $K(\mathbb{Z}_k, 1)$ has infinite cohomological dimension.

Finally, let us show that $\pi_1(N)$ contains a finite index normal abelian subgroup. Since $\pi_1(M)$ is a Bieberbach group, there exists a finite index normal subgroup $\mathbb{Z}^d \subseteq \pi_1(M)$. The image of $\mathbb{Z}^d$ in $\pi_1(N)$ is a finitely generated normal torsion free abelian group A. Since the map $\pi_1(M)/\mathbb{Z}^d \to \pi_1(N)/A$ is surjective, A has finite index in $\pi_1(N)$. Therefore $\pi_1(N)$ is a Bieberbach group, and therefore N is homeomorphic to a Bieberbach manifold. □

Theorem 3.6 can be applied to fibration (2.1) to obtain the following corollaries.

Corollary 3.7. *If $(M, \mathcal{F})$ is a singular Riemannian foliation, and a regular leaf is homeomorphic to a flat manifold, then $(M, \mathcal{F})$ is a B-foliation.*

Corollary 3.8. *The infinitesimal foliations of a B-foliation are again B-foliations.*

3.2. **Proof of Theorem A.** We now prove that the regular leaves of a B-foliation $(M, \mathcal{F})$ on a simply connected manifold are homeomorphic to tori.

Observe that the fundamental group of M is unchanged if we discard all the singular strata of codimension at least 3. Therefore we can reduce the problem to the case of M being a union

$$M = M_{reg} \cup \bigcup_{i=1}^{r} \Sigma_i,$$

where all the singular leaves are circle bundles over the regular ones. Let L_i be the regular leaf of Σ_i, and let

$$0 \to \mathbb{Z} \to \pi_1(L) \to \pi_1(L_i) \to 1$$

be the short exact sequence of the circle bundle $\mathsf{p}_i : L \to L_i$. Denote by k_i the generator of the kernel of p_{i*}, and let $K \subseteq \pi_1(L)$ be the group generated by the elements k_i.

By recent work of Goertsches, Lytchak and Toeben, there is a commutative diagram

$$\begin{tikzcd} L \arrow[r, "\hat{\iota}"] \arrow[d, equal] & \hat{M}_{reg} \arrow[r] \arrow[d, "f"] & B \arrow[d] \\ L \arrow[r, "\iota"] & M_{reg} \arrow[r] & M_{reg}/\mathcal{F} \end{tikzcd} \tag{3.1}$$

where the first row is a fibration, L is a regular leaf of $\mathcal{F}$, f is a homotopy equivalence, and B is the classifying space of the orbifold $M_{reg}/\mathcal{F}$. This commutative diagram can be obtained by first taking the Molino bundle (cf. [32, Section 5.2])

$$\begin{tikzcd} E^1_T \arrow[r, "\pi^1_T"] \arrow[d] & W \arrow[d] \\ M_{reg} \arrow[r] & M_{reg}/\mathcal{F} \end{tikzcd} \tag{3.2}$$

where E^1_T is the *transverse frame bundle*, i.e. the bundle of frames orthogonal to the leaves, and W is the frame bundle of $M_{reg}/\mathcal{F}$. Recall that W is always a manifold. The map $\pi^1_T : E^1_T \to W$ is equivariant under the action of $\mathrm{O}(q)$, where q is the codimension $\mathcal{F}$. Applying the Borel construction to E^1_T and W one obtains an induced map

$$^*\pi^1_T : E^1_T \times_{\mathrm{O}(q)} E\mathrm{O}(q) \to W \times_{\mathrm{O}(q)} E\mathrm{O}(q).$$

We then let

$$\begin{aligned} \hat{M}_{reg} &= E^1_T \times_{\mathrm{O}(q)} E\mathrm{O}(q), \\ B &= W \times_{\mathrm{O}(q)} E\mathrm{O}(q), \end{aligned}$$

and the diagram (3.1) follows from the naturally of the Molino bundle (3.2).

Let H be the image of the boundary map $\pi_2(B) \to \pi_1(L)$ in the homotopy exact sequence of $L \to \hat{M}_{reg} \to B$. There is an exact sequence

$$0 \to H \to \pi_1(L) \xrightarrow{\hat{\iota}} \pi_1(\hat{M}_{reg}) \to \pi_1(B) \to 1. \tag{3.3}$$

We will prove the result by showing that $\pi_1(L)$ is abelian. To do so, we will proceed in two steps:

Step 1: $K \subseteq \pi_1(L)$ maps surjectively onto $\pi_1(M_{reg})$ under $\hat{\iota}_*$. In particular, B is simply connected.

Step 2: K is abelian, and $H \subseteq \pi_1(L)$ is central.

By the first step, the exact sequence (3.3) becomes

$$0 \to H \to \pi_1(L) \xrightarrow{\hat{\iota}} \pi_1(\hat{M}_{reg}) \to 1.$$

By the second step, $\pi_1(\hat{M}_{reg})$ is abelian and $\pi_1(L)$ is a central extension of two abelian groups, and therefore abelian.

Proof of Step 1. Define the sets $M_j \subseteq M$ by

$$M_j = M_{reg} \cup \bigcup_{i=1}^{j} \Sigma_j,$$

and let $G_j = \pi_1(M_j)$. In particular, $G_0 = \pi_1(M_{reg})$ and $G_r = \pi_1(M) = \{1\}$. Let $g_i = \iota_* k_i$. We will prove by induction that $G_j \simeq G_0/\langle g_1, \dots g_k\rangle$, which will prove the first step.

We compute the groups G_i inductively via the van Kampen theorem. Taking an open cover $\{U, V\}$ of M_i, where $U = M_{i-1}$ and $V = M_{reg} \cup \Sigma_i$, we obtain:

$$\begin{tikzcd}
G_{i-1} \arrow[r] & G_i \\
\pi_1(\nu\Sigma_i) \arrow[u, "J_{i*}"] \arrow[r, "Q_{i*}"] & \pi_1(\Sigma_i) \arrow[u] \\
\pi_1(L) \arrow[u, "\overline{\lambda}_{i*}"] \arrow[r, "\mathsf{p}_{i*}"] & \pi_1(L_i) \arrow[u, "\lambda_{i*}"']
\end{tikzcd} \tag{3.4}$$

The upper square corresponds to the van Kampen Theorem, while the lower one comes from the commutative diagram of fibrations

$$\begin{tikzcd}
\mathbb{S}^1 \arrow[r] & \nu\Sigma_i \arrow[r, "Q_i"] & \Sigma_i \\
\mathbb{S}^1 \arrow[r] & L \arrow[u, "\overline{\lambda}_i"] \arrow[r, "\mathsf{p}_i"] & L_i \arrow[u, "\lambda_i"']
\end{tikzcd} \tag{3.5}$$

that makes p_i the restriction of Q_i to L_i.

Let $\overline{g}_i$ denote a generator of the kernel of Q_{i*}. By the van Kampen Theorem, $G_i \simeq G_{i-1}/\langle J_{i*}\overline{g}_i\rangle$.

Assume, by induction, that $G_{i-1} \simeq G_0/\langle g_1, \dots g_{i-1}\rangle$ so that the following diagram commutes:

$$\begin{tikzcd}
G_0 \arrow[r, "(M_0 \subseteq M_{i-1})_*"] \arrow[dr] & G_{i-1} \arrow[d, "\simeq"] \\
& G_0/\langle g_1, \dots g_{i-1}\rangle
\end{tikzcd} \tag{3.6}$$

Diagram (3.6) above fits in the commutative diagram

$$\begin{tikzcd}
 & \pi_1(L) \arrow[dl, "\iota_*"'] \arrow[d, "J_{i*}\overline{\lambda}_{i*}"] & \\
G_0 \arrow[r] \arrow[dr] & G_{i-1} \arrow[r] \arrow[d, "\simeq"] & G_i \simeq G_{i-1}/\langle J_{i*}\overline{g}_i\rangle \\
 & G_0/\langle g_1, \dots g_{i-1}\rangle \arrow[ur] &
\end{tikzcd}$$

Now, $\overline{g}_i = \overline{\lambda}_{i*}k_i$, and $J_{i*}\overline{g_i} = J_{i*}\overline{\lambda}_{i*}k_i$ corresponds to the image of g_i in $G_0/\langle g_1, \dots g_{i-1}\rangle$. Therefore,

$$G_i \simeq G_{i-1}/\langle J_{i*}\overline{g}_i\rangle \simeq G_0/\langle g_1, \dots g_i\rangle$$

and the induction step is proved.

Proof of Step 2. Suppose that $k_i, k_j \in K$ do not commute or, equivalently, that

$$k_i k_j k_i^{-1} \neq k_j. \tag{3.7}$$

If we consider the circle bundle $\mathsf{p}_j : L \to L_j$, equation (3.7) can be restated as saying that p_j is not orientable along a curve representing the class k_i. Therefore, we must have

$$k_i k_j k_i^{-1} = k_j^{-1}$$

or, equivalently,

$$[k_i, k_j] = k_i^{-2}.$$

By exchanging the roles of k_i and k_j, we similarly obtain

$$[k_i, k_j] = k_i^{-2}.$$

Thus

$$k_i^{-2} = k_j^{-2}.$$

If k_i and k_j do not commute, the same is true for k_i and k_j^{-1}. Repeating the same argument as before, we obtain

$$k_i^{-2} = [k_i, k_j^{-1}] = k_j^2.$$

It follows that $k_j^2 = k_i^{-2} = k_j^{-2}$, contradicting the fact that $\pi_1(L)$ is torsion free. $\square$

4. EULER CHARACTERISTIC OF B-FOLIATIONS

A well-known theorem of Kobayashi states that, if M is a smooth manifold with a smooth effective torus action, then $\chi(M)$, the Euler characteristic of M, is the Euler characteristic of the fixed point set of the torus action (cf. [30]). The goal of this section is to prove Theorem B, which is a generalization of Kobayashi's theorem to B-foliations.

We will first assume the existence of a good open cover for the leaf space $M/\mathcal{F}$ of a singular Riemannian foliation $(M,\mathcal{F})$, i.e. an open cover $\{\,U_\alpha\,\}_\alpha$ of $M/\mathcal{F}$ such that every finite intersection $U_{\alpha_1}\cap\ldots\cap U_{\alpha_k}$ is contractible. We present a proof of this assertion in the following subsection (cf. Theorem 4.1 below).

4.1. **Proof of Theorem B.** Let W be a small tubular neighborhood $B_\epsilon(\Sigma_0)$ of Σ_0 and let $V = M\setminus B_{\epsilon/2}(\Sigma_0)$. As $\{\,W,V\,\}$ is an open cover of M, we have

$$\begin{aligned}\chi(M) &= \chi(W)+\chi(V)-\chi(W\cap V)\\ &= \chi(\Sigma_0)+\chi(V)-\chi(W\cap V).\end{aligned}$$

Observe that V and $W\cap V$ are saturated submanifolds without 0-dimensional leaves. We will now show that $\chi(V\cap W)=\chi(V)=0$, which proves the theorem. Notice that $V\cap W$ retracts to $M'=\partial B_{\epsilon/2}(\Sigma_0)$, which is a compact saturated submanifold of M. In particular, it follows from Theorem 4.1 below that $M'/\mathcal{F}|_{M'}$ admits a finite good open cover $\{U_1,\ldots,U_k\}$. Therefore,

$$\chi(M')=\sum_i \chi(U_i)-\sum_{i,j}\chi(U_i\cap U_j)+\ldots,$$

where the sum is finite. Observe that every finite intersection retracts to a leaf. Since these leaves are homeomorphic to nontrivial Bieberbach manifolds, their Euler characteristic is 0. In particular, $\chi(M')=\chi(W\cap V)=0$. We will now show that $\chi(V)=0$. Observe first that $\overline{V}$, the closure of V, has boundary M'. The double $M''=\overline{V}\cup_{M'}\overline{V}$ is a compact manifold which admits a B-foliation without 0-dimensional leaves and, as before, $\chi(M'')=0$. On the other hand, $\chi(M'')=2\chi(V)-\chi(M')$, so $\chi(V)=0$. This concludes the proof of Theorem B. □

4.2. **Good covers of leaf spaces.** We now prove the existence of a good open cover for the leaf space of a singular Riemannian foliation. This has also been proven by Wolak [49]. We include a complete proof here for the benefit of the reader. We closely follow, with due changes, the proof of the existence of good open covers in Riemannian manifolds.

Theorem 4.1. *The leaf space $M/\mathcal{F}$ of a singular Riemannian foliation $(M,\mathcal{F})$ admits a good open cover.*

Proof. Showing the existence of a good open cover of $M/\mathcal{F}$ is equivalent to proving the existence of a saturated open cover $\{\overline{U}_\alpha\}_\alpha$ of M such that every finite intersection $\overline{U}_{\alpha_1}\cap\ldots\cap\overline{U}_{\alpha_k}$ contracts to one of its leaves. We start with two preliminary lemmas and a proposition; we then prove the existence of good open covers in Proposition 4.5.

Lemma 4.2. *For every $L \in \mathcal{F}$, there exist ϵ, $\delta > 0$ such that, for every other leaf $L' \subseteq \Sigma_L \cap B_\epsilon(L)$, the neighborhood $B_\delta(L')$ is a normal tubular neighborhood of L', and $B_\epsilon(L) \subseteq B_\delta(L')$.*

Proof of Lemma 4.2. Let L_x be the leaf corresponding to the point $x \in \Sigma_L/\mathcal{F}_{\Sigma_L} \subseteq M/\mathcal{F}$ and define $F : \Sigma_L/\mathcal{F}_{\Sigma_L} \to \mathbb{R}$ by

$$F(x) = \max\{\, r \mid B_r(L_x) \text{ is a normal tubular neighborhood of } L_x \,\}.$$

Let $\overline{F}(r) := \inf\{F|_{B_r(L)}\}$. Since F is continuous, $\overline{F}$ is continuous as well. Moreover, $\overline{F}$ is decreasing and $\lim_{r\to 0} \overline{F}(r) = F(L)$. Therefore, there exists $\epsilon > 0$ such that $\overline{F}(\epsilon) > \epsilon$ and, if δ is any number in $(\epsilon, \overline{F}(\epsilon))$, then ϵ and δ satisfy the conclusions of the lemma. □

Lemma 4.3. *For every $L \in \mathcal{F}$, there exists $\eta > 0$ such that, for every $0 < r < \eta$ and $L' \in S_r(L)$, any horizontal geodesic starting at L' tangent to $S_r(L')$ leaves $B_r(L)$.*

Proof of Lemma 4.3. Consider the subset $D \subseteq T^1M$ given by horizontal unit vectors, with projection $\pi_D : D \to M$. Fix $\epsilon > 0$ and define the function

$$G : D \cap B_\epsilon(L) \to \mathbb{R}$$

by

$$G(p, v_p) = \frac{d^2}{dt^2}\Big|_{t=0} \operatorname{dist}(\gamma_{v_p}, L),$$

where γ_{v_p} is the horizontal geodesic starting at (p, v_p). Since $G|_{\pi_D^{-1}(L)} > 0$, there is a neighborhood W of $\pi_D^{-1}(L)$ such that $G|_W > 0$. By shrinking W if necessary, we can suppose that

$$W = \left\{\, (p, v_p) \mid p \in B_\eta(L), v_p \in \pi_D^{-1}(L_p) \,\right\}.$$

Any η in the definition of W will satisfy the conclusion of the lemma. □

Proposition 4.4. *For every $L \in \mathcal{F}$, there exists $\rho > 0$ such that for every pair of leaves L_1, $L_2 \subseteq B_\rho(L)$, and for every minimizing geodesic γ from L_1 to L_2, the interior of γ is contained in $B_\rho(L)$.*

Proof of Proposition 4.4. Take η as in Lemma 4.3 and fix $\rho < \eta/2$. We will show that $B_\rho(L)$ satisfies the conclusion of the proposition. Observe that a geodesic γ between two leaves $L_1, L_2 \subseteq B_\rho(L)$ has length $l(\gamma) < 2\rho$. Moreover, for each t, we have

$$\begin{array}{rcl} \operatorname{dist}(\gamma(t), L) & \leq & l(\gamma)/2 + \rho \\ & < & 2\rho \\ & < & \eta. \end{array}$$

Therefore, $\mathrm{Im}(\gamma) \subseteq B_\eta(L)$. If γ were not contained in $B_\rho(L)$, the function $\mathrm{dist}(\gamma(t), L)$ would have a maximum for some t in the interior, contradicting Lemma 4.3. $\square$

We are now ready to prove the existence of a good open cover for leaf spaces. For every $L \in \mathcal{F}$, let ρ_L be as in Proposition 4.4, let ϵ_L be as in Lemma 4.2, and take $R_L < \min\{\, \rho_L,\ \epsilon_L \,\}$.

Proposition 4.5. *The open cover $\{B_{R_L}(L)\}_{L\in\mathcal{F}}$ projects to a good open cover of $M/\mathcal{F}$.*

Proof of Proposition 4.5. Let $U = \bigcap_{i=1}^{N} B_{R_i}(L_i)$. Since $R_i < \epsilon_i$, we know by the homothetic transformation lemma that the stratification of $\mathcal{F}$ restricted to $B_{R_i}(L_i)$ consists of a chain and, in particular, there is a unique minimal stratum Σ_i. Therefore, the stratification in U also admits a minimal stratum, which we will call Σ_0. Suppose that $L_1 \in \Sigma_0$. Then, given a leaf L in $U \cap \Sigma_0$, by Lemma 4.2 there is a tubular neighborhood $B_\delta(L) \supseteq D_{R_1}(L_1) \supseteq U$, and in particular for every L' in U there is a “unique” geodesic γ from L to L', contained in $B_\delta(L)$. Here by “unique” we mean that for each $p \in L$, there exists a unique minimizing geodesic from L to L' starting at p. By Proposition 4.4, γ is contained in every $B_{R_i}(L_i)$, and therefore $Im(\gamma) \subseteq U$. In particular, it is possible to retract U to L along these horizontal geodesics. In other words, it is possible to retract $\pi(U)$ to the point $\pi(L)$, where $\pi : M \to M/\mathcal{F}$ is the canonical projection. Therefore $\pi(U) = \bigcap_{i=1}^{N} \pi(B_{R_i}(L_i))$ is contractible. $\square$

$\square$

Remark 4.6. Alexander Lytchak has pointed out to us that one may prove an analog of Theorem 4.1 for Alexandrov spaces proceeding along the lines of the proof we have presented here. In this more general setting the role of tubular neighborhoods and geodesics is played by convex neighborhoods and flow lines of strictly concave functions, whose existence has been proved by Perelman and Petrunin [41].

5. B-foliations of codimension 1 on simply connected manifolds

In this section we classify up to foliated diffeomorphism all possible codimension 1 B-foliations on simply connected manifolds (see Theorem 5.2 below). We first prove the following lemma.

Lemma 5.1. *Let M^{n+1} be a compact, simply connected $(n+1)$-manifold. If $(M^{n+1}, \mathcal{F}^n)$ is a codimension 1 B-foliation, then the foliation must be singular.*

Proof. Suppose that $\mathcal{F}^n$ is a regular foliation. Since M^{n+1} is simply connected, it follows from work of Molino [32] that $\mathcal{F}^n$ must be a simple foliation, i.e. it is given by the fibers of a Riemannian submersion. Hence there is a fibration $L^n \to M^{n+1} \to \mathbb{S}^1$. Since M^{n+1} is simply connected, the long exact sequence in homotopy for the fibration yields a contradiction. $\square$

Theorem 5.2. *The only codimension* 1 *B-foliations on compact simply connected manifolds are the homogeneous foliations* $(\mathbb{S}^2, \mathbb{S}^1)$ *and* $(\mathbb{S}^3, T^2)$.

Proof. Let M be a compact, simply connected manifold and $(M, \mathcal{F})$ a codimension 1 B-foliation of M. By Lemma 5.1, $\mathcal{F}$ is singular. Therefore, the leaf space M^* is homeomorphic to a closed interval $[-1, 1]$. Let L_t be the leaf corresponding to $t \in [-1, 1]$ and observe that the manifold M decomposes as the union of two disk bundles over $L_{\pm 1}$. If L_0 is regular, there are submersions $\pi_t : L_0 \to L_t$. By [31, Theorem 1.6], the leaves $L_{\pm 1}$ are singular, and there are submersions $\mathbb{S}^r \to L_0 \to L_{\pm 1}$. Since L_0 and $L_{\pm 1}$ are diffeomorphic to Bieberbach manifolds, we must have that $\mathbb{S}^r = \mathbb{S}^1$. By [20, Table 1.4], the inclusion map $L_0 \hookrightarrow M$ is homotopic to a fibration $F \to L_0 \to M$, with $\pi_1(F)$ isomorphic to one of Q, $\mathbb{Z} \oplus \mathbb{Z}$, or $\mathbb{Z} \oplus \mathbb{Z}_2$, where Q denotes the order 8 subgroup $\{ \pm 1, \pm i, \pm j, \pm k \}$ of the unit quaternions $\mathbb{S}^3 \subset \mathbb{H}$. The long exact sequence in homotopy of the fibration $F \to L_0 \to M$ implies that $\pi_1(F) \to \pi_1(L_0) \to 0$. Therefore, $\pi_1(L_0)$ can contain at most a $\mathbb{Z}^2$ subgroup. It follows from Bieberbach's theorem that L_0 is at most 2-dimensional, and hence M has dimension at most 3. Since M is simply connected, it must be diffeomorphic to $\mathbb{S}^2$ or $\mathbb{S}^3$.

In the case of $\mathbb{S}^2$, we know from Theorem B, or by inspection of the leaf space, that the set of zero dimensional leaves consists of 2 points p_1, p_2, and the regular leaves are the distance spheres from these two points. The regular part is foliated diffeomorphic to $((0, 1) \times \mathbb{S}^1, \mathbb{S}^1)$, where the $\mathbb{S}^1$-action fixes the first factor. In particular, there exists a unit vector field X defined on the regular part of $\mathbb{S}^2$, spanning the foliation there. We can now define a new vector field $X'_p = \ell(L_p) \cdot X_p$, where $\ell(L_p)$ denotes the length of L_p, and extend X' to the singular part by 0. One can verify that this vector field is smooth (see, for example, the proof of Proposition 8.5 below) and the foliation in $\mathbb{S}^2$ is given by the orbits of a smooth circle action.

Suppose now that M is diffeomorphic to $\mathbb{S}^3$. In this case, the regular leaves are 2-dimensional, the singular leaves $L_{\pm 1}$ are 1-dimensional, hence they are circles, and the regular leaves fiber over $L_{\pm 1}$ with fiber a circle. By Theorem A, the regular leaves are diffeomorphic to a 2-torus. We have that $\mathbb{S}^3$ can be described as a double disk bundle

$$M = D_- \cup_\phi D_+,$$

where $D_\pm$ is a disk bundle over $L_{\pm 1}$ and $\phi : \partial D_+ \to \partial D_-$ is a diffeomorphism. Moreover, the foliation $\mathcal{F}$ consists of the distance tubes to the zero section (with respect to some Euclidean structure on the disk bundles). Notice that $D_\pm$ are solid tori and, $\partial D_\pm$ are tori.

We will now show that the foliation is homogeneous. Consider cylindrical coordinates (z, θ, r) on D_+ and (z', θ', r') on D_-. In this way, we can consider the diffeomorphism $\phi : \partial D_+ \to \partial D_-$ as an element of $\mathrm{Diff}(T^2)$. All such diffeomorphisms are isotopic to a map that is linear with respect to the chosen cylindrical coordinates. Since the isotopy class of the gluing preserves the diffeomorphism type of M, we can assume, without loss of generality, that ϕ is a linear map given by a matrix $A \in \mathrm{SL}(2, \mathbb{Z})$. In our case, since M is $\mathbb{S}^3$, we may take

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Moreover, we can modify the Riemannian metric radially, so that it becomes a product near the boundary and the foliation remains Riemannian. One can explicitly produce a smooth T^2 action that gives rise to $\mathcal{F}$ as follows. Given $(s,t) \in T^2$, define an action of (s,t) on $D_\pm$ by

$$\begin{aligned} (s,t) \star (z, \theta, r) &= (z+s, \theta+t, r), \\ (s,t) \star (z', \theta', r') &= (z'+t, \theta'+s, r'), \end{aligned}$$

where $\phi = (\phi_1, \phi_2)$ is the gluing diffeomorphism $\phi : \partial D_+ \to \partial D_-$. By the classification of cohomogeneity 1 torus actions on lens spaces (see, for example [34]), these actions are all equivariantly diffeomorphic to orthogonal actions. ☐

6. B-FOLIATIONS OF CODIMENSION 2 ON SIMPLY CONNECTED MANIFOLDS

In this section we prove Theorem C, which describes the leaf space structure of codimension 2 B-foliations on compact, simply connected manifolds. This theorem generalizes to B-foliations results on the structure of the orbit space of cohomogeneity two smooth torus actions on simply connected manifolds (cf. [34, 33, 28]).

6.1. **Regular B-foliations of codimension 2.**

We first consider the case where the codimension 2 B-foliation is regular.

Proposition 6.1. *Let $(M, \mathcal{F})$ be a B-foliation of codimension 2 on a compact simply connected manifold M. If $\mathcal{F}$ is regular, then M is diffeomorphic to $\mathbb{S}^3$ and the generic leaf is diffeomorphic to $\mathbb{S}^1$.*

Proof. Since $\mathcal{F}$ is regular, TM splits as a sum of bundles $TM = T\mathcal{F} \oplus T\mathcal{F}^\perp$, where $T\mathcal{F}$ is the vertical tangent bundle and $T\mathcal{F}^\perp$ is the horizontal tangent bundle. Let F_H be the frame bundle of $T\mathcal{F}^\perp$. Since $\mathcal{F}$ is of codimension 2, we have a fibration $\mathbb{S}^1 \to F_H \to M$ and its long exact homotopy sequence yields the sequence

$$\mathbb{Z} \longrightarrow \pi_1(F_H) \longrightarrow 0.$$

It follows that $\pi_1(F_H)$ is isomorphic to 0, $\mathbb{Z}$ or $\mathbb{Z}_k$ for some integer $k > 2$.

Consider now the *lifted foliation* $\mathcal{F}'$ on F_H. By Molino [32], the leaves of $\mathcal{F}'$ are finite covers of the leaves in $\mathcal{F}$, and are given by the preimages of a submersion $F_H \longrightarrow W$ on a manifold W, so W must be 3-dimensional.

Let L be the generic leaf of $(M, \mathcal{F})$. Since $(M, \mathcal{F})$ is a B-foliation, $\pi_k(L) = 0$ for $k \geq 2$. There is a fibration $L \to F_H \to W$ and the end of its long exact homotopy sequence reads

$$0 \to \pi_2(F_H) \to \pi_2(W) \xrightarrow{f} \pi_1(L) \xrightarrow{g} \pi_1(F_H) \xrightarrow{h} \pi_1(W) \to 1. \tag{6.1}$$

Now, if $\pi_1(F_H)$ is finite, then $\pi_1(W)$ is also finite, so W is finitely covered by $\mathbb{S}^3$. In particular, $\pi_2(W) = 0$ and there is an injection

$$0 \to \pi_1(L) \to \pi_1(F_H)$$

with $\pi_1(F_H)$ finite, which contradicts the Bieberbach theorem.

It follows that $\pi_1(F_H) = \mathbb{Z}$, and $\pi_1(W)$ must be isomorphic to 0, $\mathbb{Z}_k$, for $k > 2$, or to $\mathbb{Z}$. If $\pi_1(W) = \mathbb{Z}$, then the map $h : \pi_1(F_H) \to \pi_1(W)$ is an isomorphism and $W \simeq \mathbb{S}^1 \times \mathbb{S}^2$. In particular $\pi_2(W) = \mathbb{Z}$ and, from the long exact sequence (6.1), we get the short exact sequence

$$0 \to \pi_2(F_H) \to \mathbb{Z} \xrightarrow{f} \pi_1(L) \longrightarrow 0.$$

Since L is a Bieberbach manifold, we must have that $\pi_1(L) = \mathbb{Z}$, $L \simeq \mathbb{S}^1$, f is injective and $\pi_2(M) = 0$. Since L is 1-dimensional, F_H has dimension 4 and M has dimension 3, so the only possibility is $(M, \mathcal{F}) = (\mathbb{S}^3, \mathbb{S}^1)$.

If $\pi_1(W) = 0$ or $\mathbb{Z}_k$, then W is finitely covered by $\mathbb{S}^3$ and $\pi_2(W) = 0$. The homotopy sequence then becomes

$$0 \to \pi_2(F_H) \to 0 \xrightarrow{f} \pi_1(L) \xrightarrow{g} \mathbb{Z} \xrightarrow{h} \pi_1(W) \to 1.$$

As before, L has to be $\mathbb{S}^1$, M is 3-dimensional and $(M, \mathcal{F}) = (\mathbb{S}^3, \mathbb{S}^1)$. □

Remark 6.2. By definition, a codimension 2 regular B-foliation on $\mathbb{S}^3$ corresponds to a Seifert fibration on $\mathbb{S}^3$. It follows from the classification of Seifert manifolds that every possible B-foliation of $\mathbb{S}^3$ can be realized by a smooth circle action (cf. [36]).

6.2. **Singular B-foliations of codimension 2.** We now consider the case where the codimension 2 B-foliation is singular.

Let $(M, \mathcal{F})$ be a codimension 2 B-foliation. By [31, Theorem 1.6], if the foliation is not regular, then there are no exceptional leaves. Moreover, the leaf space M^* is homeomorphic to a 2-dimensional orbifold B. By [18, Lemma 3.5], the topological interior $B \setminus \partial B$ of B is simply connected, and therefore is homeomorphic to an open disk. The boundary of B consists of the union of geodesic arcs. The points in the interior of these arcs correspond to leaves which we will call by *least singular leaves*, while the vertices of the leaf space, i.e. the points where two geodesic arcs in the boundary meet, correspond to leaves which we will call *most singular leaves* (see Figure 1).

Let L be a singular leaf and fix $p \in L$. By Corollary 3.8, the infinitesimal foliation $(\mathbb{S}^r_p, \mathcal{F}_p)$ is a codimension one B-foliation, whose quotient is a closed interval. By Theorem 5.2, the infinitesimal foliation $(\mathbb{S}^r_p, \mathcal{F}_p)$ must be one of the homogeneous foliations $(\mathbb{S}^2, \mathbb{S}^1)$ or $(\mathbb{S}^3, T^2)$. Since $\mathbb{S}^r_p$ is a round unit sphere, it follows from the main theorem in [43] that the foliation is isometric to one of the homogeneous foliations $(\mathbb{S}^2, \mathbb{S}^1)$ or $(\mathbb{S}^3, T^2)$ induced by orthogonal actions on round unit spheres. In the first case, the leaf space of the homogeneous foliation is isometric to a closed interval of length π and corresponds to the infinitesimal foliation of a least singular leaf. In the



FIGURE 1. Example of a leaf space of a codimension 2 B-foliation.

second case, the leaf space of the homogeneous foliation is isometric to a closed interval of length $\pi/2$ and corresponds to the infinitesimal foliation of a most singular leaf.

Since there are no exceptional leaves, by Remark 2.3, the holonomy action is trivial, and hence the space of directions at any point $p^* \in M^*$ is isometric to the leaf space of the infinitesimal foliation at p^*. In particular, the angle between geodesic arcs meeting at vertices of M^* is $\pi/2$ and the fibration 2.1 yields the desired metric fibrations in part (2) of Theorem C.

The following proposition finishes the proof of Theorem C.

Proposition 6.3. *Let $(M^{n+2}, \mathcal{F})$ be a codimension 2 B-foliation of a compact, simply connected $(n+2)$-manifold. Then the leaf space M^* hast at least n edges.*

Proof. Observe that M remains simply connected if we discard all the singular strata of codimension strictly greater than 2, i.e. all those leaves corresponding to the corners in the leaf space M^*. As in the first step in the proof of Theorem A, we consider

$$M = M_{reg} \cup \bigcup_{j=1}^{r} \Sigma_j,$$

where the strata Σ_j project to the edges of M^*. We let

$$M_i = M_{reg} \cup \bigcup_{j=1}^{i} \Sigma_j,$$

and $G_i = \pi_1(M_i)$.

We will show, by induction on i, that G_i contains a copy of $\mathbb{Z}^{r_i}$ of finite index, where $r_i \geq n - i$. This will prove the proposition, since $G_r = \{1\}$ and therefore it would follow that $n - r \leq 0$.

Since $G_0 = \pi_1(L) = \mathbb{Z}^n$, the initial step of the induction holds. Suppose now that the induction hypothesis holds for G_{i-1}.

Again, from the first step in the proof of Theorem A, $G_i = G_0/\langle g_1, \dots g_i\rangle$ and therefore there are exact sequences

$$0 \to \langle g_i\rangle \to G_{i-1} \to G_i \to 1,$$

where $\langle g_i\rangle$ either generates a $\mathbb{Z}_k$ or a $\mathbb{Z}$. In either case, $\mathbb{Z}^{r_{i-1}} \subseteq G_{i-1}$ projects to a finitely generated abelian group $A \subseteq G_i$ of finite index, and A_F, the free part of A, has also finite index in G_i. Observe that A is isomorphic to $\mathbb{Z}^{r_{i-1}}/(\langle g_i\rangle \cap \mathbb{Z}^{r_{i-1}})$. The subgroup $\langle g_i\rangle \cap \mathbb{Z}^{r_{i-1}}$ is isomorphic to $\mathbb{Z}$, if $\langle g_i\rangle = \mathbb{Z}$ and $\langle g_i\rangle \cap \mathbb{Z}^{r_{i-1}} \neq \emptyset$, or 0 otherwise. Therefore, A_F will be $\mathbb{Z}^{r_{i-1}}$ or $\mathbb{Z}^{r_{i-1}-1}$. Either way,

$$r_i \geq r_{i-1} - 1 \geq n - (i-1) - 1 = n - i,$$

which proves the induction step. □

7. Curvature and B-foliations of codimension 2

In this section, we apply the results in the previous sections to the study of compact, simply connected Riemannian manifolds with positive or nonnegative sectional curvature and a B-foliation of low codimension. In particular, we prove Theorems C and D, which generalize results in [17] to the case of B-foliations.

This section is motivated by the wealth of results on positively and nonnegatively compact Riemannian manifolds with large effective isometric torus actions (see references in the Introduction) and our results may be cast in a similar light. More precisely, recall that the *symmetry rank* of a compact Riemannian manifold M is, by definition, the rank of the isometry group of M (cf. [21]). Therefore, M supports a "large" torus action if it has big symmetry rank. If M is n-dimensional and has positive sectional curvature, Grove and Searle [21] showed that the symmetry rank of M is bounded above by $\lfloor (n+1)/2 \rfloor$, and classified compact Riemannian manifolds of positive curvature and maximal symmetry rank. In view of these results, it seems natural to study positively and nonnegatively compact Riemannian manifolds with B-foliations of low codimension, in analogy with positively and nonnegatively curved manifolds with large symmetry rank.

The simplest examples of B-foliations on positively and nonnegatively curved manifolds, standing closest to those induced by isometric torus actions, are B-foliations whose regular leaves are flat with the induced metric. We call these foliations *Euclidean foliations* (see Definition 7.1 below). We will make some observations on these foliations in the following subsection, framing Theorems C and D in this context. We will then prove Theorem D in Subsection 7.2, and Theorem E in Subsection 7.3.

7.1. **Quasi-positive curvature and Euclidean foliations.**

We now introduce *Euclidean foliations* and the *Euclidean rank* of a Riemannian manifold. The latter invariant is a generalization of the symmetry rank of a Riemannian manifold (cf. [21]).

Definition 7.1 (Euclidean foliation)**.** A singular Riemannian foliation $(M, \mathcal{F})$ is *Euclidean* if it is a B-foliation and the regular leaves are flat with the induced metric.

Definition 7.2 (Euclidean rank)**.** The *Euclidean rank* of a Riemannian manifold (M, g) is defined by

$$\operatorname{Erank}(M, g) = \max\{\, \dim \mathcal{F} \mid (M, \mathcal{F}) \text{ is a Euclidean foliation}\,\}.$$

Remark 7.3. In the definition of the Euclidean rank, the Riemannian metric is fixed and then we maximize over all the possible Euclidean singular Riemannian foliations compatible with this metric. If one in turn maximizes the Euclidean rank over all Riemannian metrics of a smooth manifold M, one gets an invariant of M akin to the degree of symmetry of a smooth manifold.

Recall that a Riemannian manifold (M, g) is said to have *quasi-positive curvature* if (M, g) has nonnegative (sectional) curvature and a point with strictly positive curvature. The following bound for the Euclidean rank of a quasi-positively curved manifold follows from Otsuki's lemma [9, Lemma 3.3, p. 224] (cf. also an argument due to Burkhard Wilking found in [27, 16]).

Theorem 7.4. *If (M, g) is a quasi-positively curved Riemannian n-manifold, then*

$$\operatorname{Erank}(M, g) \leq \lfloor (n+1)/2 \rfloor.$$

In view of Theorem 7.4, one would like to classify, up to diffeomorphism, positively curved manifolds of maximal Euclidean rank, thus extending Grove and Searle's classification of positively curved manifolds with maximal symmetry rank (cf. [21]). A difficulty arises in the foliated case, though, since an analogue of Berger's lemma is lacking. This result implies that a torus action of maximal rank on a positively curved manifold contains a circle subgroup with a codimension 2 fixed point set, which is a crucial step in Grove and Searle's classification. Nevertheless, one can obtain a topological classification in dimensions 4 and 5, where the leaf space is of dimension 2 and has a relatively simple structure. In this case, one may dispense with the assumption that the foliation is Euclidean, and it suffices to assume the existence of codimension 2 B-foliation. We obtain in this way Theorem D.

One may also relax the curvature assumptions and consider the Euclidean rank of compact, simply connected manifolds with nonnegative curvature. Let M^{n+2} be a simply connected, compact Riemannian n-manifold with a codimension 2 B-foliation and nonnegative curvature. By Theorem 6.3, the leaf space M^* has at least n vertices. On the other hand, by a comparison argument (see the proof of Theorem D below), we must have $n \leq 4$ and M must be at most 6-dimensional. It follows that, in dimensions $n \leq 9$, the Euclidean rank of a compact, simply connected Riemannian n-manifold of nonnegative curvature is bounded above by $\lfloor 2n/3\rfloor$. The bound on the Euclidean rank in dimensions 4, 5 and 6 is sharp, since there exist nonnegatively curved simply connected manifolds with cohomogeneity two isometric torus actions, for example, $\mathbb{S}^2\times\mathbb{S}^2$, $\mathbb{S}^3\times\mathbb{S}^2$ and $\mathbb{S}^3\times\mathbb{S}^3$ with their usual nonnegatively curved product metrics.

The classification of compact, simply connected Riemannian 4- and 5-manifolds with nonnegative curvature and a codimension 2 B-foliation is given by Theorem E. In particular, this theorem yields a topological classification of compact, simply connected Riemannian 4- and 5-manifolds of nonnegative curvature and maximal Euclidean rank, generalizing results in [17].

We now present the proofs of Theorems D and E.

7.2. **Proof of Theorem D.** We prove assertions (1) and (2) separately.

Proof of assertion (1). Suppose $n=4$. Then the leaf space is a 2-dimensional Riemannian manifold of nonnegative curvature, with polyhedral boundary, and with positive curvature in an open subset. By Theorem 6.3, the leaf space M^* has at least two vertices p_1^* and p_2^* corresponding to 0-dimensional leaves and, by Theorem A, the regular leaves are 2-tori.

By Theorem C, the angles at the vertices of the leaf space M^* are equal to $\pi/2$. Suppose now that there are five vertices $p_0^*,\ldots,p_4^*$ in ∂M^*, and assume that the points p_i^*, p_{i+1}^*, $0\le i\le 4$ are adjacent, with indices considered modulo 5. Taking minimal geodesics from p_0^* to p_2^* and p_3^*, we obtain a triangulation of M^* consisting of three triangles. By Toponogov's theorem, the sum of the angles of these three triangles is bounded below by 3π. On the other hand, adding the vertex angles at the points p_i^*, $0\le i\le 4$, we get that the sum of the angles of these three triangles is bounded above by $5\pi/2$, yielding a contradiction. Therefore, ∂M^* can only have two, three or four vertices and, if ∂M^* has four vertices, M^* must be isometric to a flat rectangle. On the other hand, since M has quasi-positive curvature, M^* has positive curvature in an open subset, and it follows from the Gauss-Bonnet theorem that M^* cannot be flat. Therefore, M^* has either two or three vertices. Since vertices correspond to points in Σ_0, the 0-dimensional stratum of the foliation, we have that Σ_0 consists of either two or three points. By Theorem B, $\chi(\Sigma_0)=\chi(M)$, so $2\le\chi(M)\le 3$. It follows from work of Freedman [13] that M is homeomorphic to $\mathbb{S}^4$, when $\chi(\Sigma_0)=2$, or to $\mathbb{C}P^2$, when $\chi(\Sigma_0)=3$. To get diffeomorphism, we now use the fact that the manifolds decompose as the union of disc bundles with boundary $\mathbb{S}^3$.

Suppose first that the quotient has two vertices p_1^* and p_2^*. Let $D_1=B_\epsilon(p_1)$, and $D_2=M-D_1$. Then M can be written as a union $M=D_1\cup_\phi D_2$, where D_1, D_2 project in the quotient to sets D_1^*, D_2^* homeomorphic to metric balls around p_1^* and p_2^*, respectively. Therefore, both D_1 and D_2 are diffeomorphic to cones over $\mathbb{S}^3$. Since $\mathbb{S}^3$ has only one smooth structure, there exist diffeomorphisms $\psi_1 : D_1\to B^4$, $\psi_2 : D_2\to B^4$ with the standard

4-ball. Letting

$$\overline{\phi} := \psi_2|_{\partial D_2} \circ \phi \circ \psi_1|_{\partial D_1}^{-1} : \mathbb{S}^3 \to \mathbb{S}^3,$$

we see that M is diffeomorphic to $B^4 \cup_{\overline{\phi}} B^4$. By Hatcher's proof of the Smale conjecture [23], $\overline{\phi}$ is isotopic to $\overline{\phi}_0 \in \mathrm{O}(4)$, and therefore M is diffeomorphic to $B^4 \cup_{\overline{\phi}_0} B^4 \simeq \mathbb{S}^4$.

Suppose now that the quotient has three vertices, and let N be the preimage of an edge in M^*. The set N is a 2-dimensional submanifold of M and $\mathcal{F}$ restricts to a codimension one foliation with two singular leaves. Hence, N id diffeomorphic to $\mathbb{S}^2$. Let now $D^1 = B_\epsilon(N)$, and $D_2 = M - D_1$. Again, D_1, D_2 project to the quotient and D_2^* is homeomorphic to a metric ball around the vertex opposite to N^*. As in the case with two vertices, there is a diffeomorphism $\psi_2 : D_2 \to B^4$, so that $\partial D_2 = \partial D_1$ is diffeomorphic to $\mathbb{S}^3$. Moreover, the $\mathbb{S}^1$-bundle $\partial D_3 \to N$ is the Hopf map $\mathbb{S}^3 \to \mathbb{S}^2$ and there is a diffeomorphism ψ_1 from D_1 to a tubular neighborhood T_1 of the standard embedding $\mathbb{S}^2 = \mathbb{C}P^1 \subseteq \mathbb{C}P^2$. Letting

$$\overline{\phi} = \psi_2|_{\partial D_2} \circ \phi \circ \psi_1|_{\partial D_1}^{-1} : \mathbb{S}^3 \to \mathbb{S}^3,$$

we conclude, as in the previous case, that $\overline{\phi}$ is isotopic to $\overline{\phi}_0 \in \mathrm{O}(4)$, and M is diffeomorphic to $T_1 \cup_{\overline{\phi}} B^4 \simeq T_1 \cup_{\phi_0} B^4$, which in turn is diffeomorphic to $\mathbb{C}P^2$.

Proof of assertion (2). By the work of Barden and Smale [3, 47], it suffices to verify that $H_2(M, \mathbb{Z}) = 0$. The leaf space M^* is homeomorphic to a disc and it has at least three vertices, by Theorem 6.3. A comparison argument as in part (1) implies that there are at most three vertices in M^*. Therefore, there are exactly three vertices.

By Theorem A, the regular leaves of the foliation are 3-tori. Observe that $\pi_1(T^3)$ is generated by γ_1, γ_2, γ_3, the kernels of the maps

$$\mathsf{p}_{i\,*} : \pi_1(T^3) \to \pi_1(T^2),$$

induced by the metric projections $\mathsf{p}_i : T^3 \to T^2$ from a regular leaf onto a least singular leaf.

Take a saturated open cover $\{U_i\}_{i=1}^3$ around the most singular leaves and let L_1, L_2 and L_3 be the most singular leaves corresponding to the vertices of the leaf space. Let U_i be neighborhoods of L_i, let $U_{ij} = U_i \cap U_j$, and let $U_0 = U_1 \cap U_2 \cap U_3$. There are retractions

$$U_i \to L_{ij} = \mathbb{S}^1,$$

$$U_{ij} \to L_{ij} = T^2$$

and

$$U_0 \to L_0 = T^3.$$

Here, L_{ij} is the generic leaf in the stratum between L_i and L_j, which corresponds to the boundary arc in the leaf space between L_i and L_j. Consider now the complex

$$C_q : \oplus H^q(U_i) \to \oplus H^q(U_{ij}) \to H^q(U_0).$$

There is a (Mayer-Vietoris) spectral sequence $H^p(C_q) \Rrightarrow H^{p+q}(M, \mathbb{Z})$. We have

$$\begin{aligned} C_0 &: \mathbb{Z}^3 \to \mathbb{Z}^3 \to \mathbb{Z}; \\ C_1 &: \mathbb{Z}^3 \to \mathbb{Z}^6 \to \mathbb{Z}^3; \\ C_2 &: 0 \to \mathbb{Z}^3 \to \mathbb{Z}^3; \\ C_3 &: 0 \to 0 \to \mathbb{Z}. \end{aligned}$$

Therefore, all the differentials are zero, so the spectral sequence stabilizes immediately. Now, we have the following sets of generators:

$$\begin{aligned} \pi_1(T^3) &= \langle \gamma_1, \gamma_2, \gamma_3 \rangle; & H^1(T^3) &= \langle \hat\gamma_1, \hat\gamma_2, \hat\gamma_3 \rangle; \\ \pi_1(L_{ij}) &= \langle \gamma_i, \gamma_j \rangle; & H^1(L_{ij}) &= \langle \hat\gamma_i, \hat\gamma_j \rangle; \\ \pi_1(L_i) &= \langle \gamma_i \rangle. & H^1(L_i) &= \langle \hat\gamma_i \rangle, \end{aligned}$$

where $\hat\gamma_i$ is the dual of γ_i. Observe that the map $\pi_1(T^3) \to \pi_1(L_{ij})$ takes $\gamma_i \mapsto \gamma_i$, $\gamma_j \mapsto \gamma_j$ and $\gamma_k \mapsto 0$, so the map

$$\mathsf{p}_i^* : H^1(L_{ij}) \to H^1(T^3)$$

is given by inclusion. In C_1, the map

$$g_1 : \mathbb{Z}^6 \simeq H^1(L_{12}) \oplus H^1(L_{13}) \oplus H^1(L_{23}) \to H^1(L_0)$$

is surjective, since

$$\hat\gamma_i \in H^1(L_{ij}) \mapsto \hat\gamma_i \in H^1(L_0).$$

In C_2, the group $H^2(L_{ij}) \simeq \mathbb{Z}$ is generated by $\hat\gamma_i \smile \hat\gamma_j$ and again the map in cohomology is an isomorphism, since g_2 takes $\hat\gamma_i \smile \hat\gamma_j \in H^2(L_{ij})$ to $\hat\gamma_i \smile \hat\gamma_j \in H^2(L_0)$. Hence, $\ker(g_2) = 0$, $\operatorname{coker}(g_1) = 0$ and, by the spectral sequence, $0 = H^3(M, \mathbb{Z}) = H_2(M, \mathbb{Z})$. □

7.3. **Proof of Theorem E.** We prove parts (1) and (2) separately.

Proof of part (1). A comparison argument as in the proof of part (1) of Theorem D shows that the leaf space may have at most four vertices. On the other hand, by Theorem 6.3, the leaf space M^* has at least two vertices. Hence, by Theorem B, the Euler characteristic of M^4 is 2, 3 or 4, and it follows from the work of Freedman [13] that M^4 is homeomorphic to $\mathbb{S}^4$, or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ or $\mathbb{S}^2 \times \mathbb{S}^2$. If M^* has two or three vertices, then the leaf space structure is the same as the one in the proof of part (1) of Theorem D and it follows that M^4 is diffeomorphic to $\mathbb{S}^4$, if M^* has two vertices, or to $\mathbb{C}P^2$, if M^* has three vertices.

Proof of part (2). By Theorem 6.3, the leaf space M^* has at least three vertices. On, the other hand, since the leaf space is nonnegatively curved, it may have at most four vertices. In the case where there are exactly four vertices, M^* must be isometric to a flat rectangle. If the leaf space has exactly three vertices, then, proceeding as in the proof of Theorem D, we conclude that M is diffeomorphic to $\mathbb{S}^5$. Therefore, we can restrict our attention to the case in which M^* is isometric to a flat rectangle, which we shall denote by $[0, x_0] \times [0, y_0]$ in the remainder of the proof. In this case, to conclude that M is diffeomorphic to one of the two $\mathbb{S}^3$-bundles over $\mathbb{S}^2$, it suffices to prove that $H_2(M, \mathbb{Z}) = \mathbb{Z}$ and then appeal to the work of Barden [3] and [47]. To prove this, we apply the Mayer-Vietoris spectral sequence as in the proof of Theorem D. Keep in mind that all the regular leaves are tori by Theorem A.

Let Y_0, Y_1 be the closures of the two strata corresponding to the vertical edges of M^*. Likewise let X_0, X_1 be the closures of the strata corresponding to horizontal edges of M^*. Since the angles between edges are all $\pi/2$, the spaces X_i, Y_i are actually 3-dimensional submanifolds of M. On each of them, the foliation is again a B-foliation of codimension 1, and the quotient is just the corresponding edge.

First step: At least one among X_i, Y_i is not homeomorphic to $\mathbb{S}^2 \times \mathbb{S}^1$. We proceed by contradiction. Suppose then that all of X_0, X_1, Y_0, Y_1 are homeomorphic to $\mathbb{S}^2 \times \mathbb{S}^1$. Let L_{X_i}, L_{Y_i} denote, as before, a generic leaf in $\mathcal{F}|_{X_i}, \mathcal{F}|_{Y_i}$, let L_0 be a generic leaf and, finally, let $S^1_{(0)}, S^1_{(1)}, S^1_{(2)}, S^1_{(3)}$ be the four leaves corresponding to the four corners of the leaf space. The metric projections give rise to a homotopy commutative diagram as shown below:

$$\begin{tikzcd}
S^1_{(1)} & L_{X_1} \arrow[l] \arrow[r] & S^1_{(3)} \\
L_{Y_0} \arrow[u] \arrow[d] & L_0 \arrow[l] \arrow[r] \arrow[u] \arrow[d] & L_{Y_1} \arrow[u] \arrow[d] \\
S^1_{(0)} & L_{X_0} \arrow[l] \arrow[r] & S^1_{(2)}
\end{tikzcd} \tag{7.1}$$

Since each X_0 is homeomorphic to $\mathbb{S}^2 \times \mathbb{S}^1$, it is easy to deduce, using the van Kampen Theorem, that the kernels of the maps $\pi_1(L_{X_0}) \to \pi_1(S^1_{(0)})$ and $\pi_1(L_{X_0}) \to \pi_1(S^1_{(2)})$ coincide. The same conclusion holds for X_1, Y_0 and Y_1. Let $x_0,\ x_1,\ y_0,\ y_1 \in \pi_1(L_0) = Z^3$ denote, respectively, the kernels of the maps $\pi_1(L) \to \pi_1(L_{X_i})$, $\pi_1(L) \to \pi_1(L_{Y_i})$ for $i = 0, 1$. Since the diagram above induces a commutative diagram on the fundamental groups, one can deduce that $x_0 = x_1 + ay_0$ and $y_0 = y_1 + bx_0$, for some $a, b \in \mathbb{Z}$. It follows that the elements x_i, y_i span a $\mathbb{Z}_2 \subseteq \pi_1(L_0)$. This fact contradicts the proof of Theorem 6.3, according to which, since M is simply connected, $\pi_1(L_0)$ should be generated by the kernels $\{x_i, y_i\}$.

Second step: $H^3(M,\mathbb{Z}) = H_2(M,\mathbb{Z}) = \mathbb{Z}$. Suppose, without loss of generality, that X_0 is not homeomorphic to $\mathbb{S}^2 \times \mathbb{S}^1$. Consider a saturated open cover $\mathfrak{U} = \{U_0, U_1, U_2\}$ of M, where the open sets are neighborhoods of $X_0, S^1_{(1)}, S^1_{(3)}$, respectively. Then $U_0 \cap U_1$, $U_0 \cap U_2$, and $U_1 \cap U_2$ are neighborhoods of L_{Y_0}, L_{Y_1}, L_{X_1} respectively, and $U_0 \cap U_1 \cap U_2$ is a neighborhood of L_0. As done before, we consider the Mayer-Vietoris spectral sequence $\check{H}^p(\mathfrak{U}, \mathcal{H}^q(\mathbb{Z})) \Rrightarrow H^{p+q}(M,\mathbb{Z})$, where the Čech cohomology is the cohomology of the following complexes:

$$\begin{aligned}
C_3 &: \bigoplus_i H^3(U_i) \to \bigoplus_{i,j} H^3(U_{ij}) \to H^3(U_{012});\\
C_2 &: \bigoplus_i H^2(U_i) \to \bigoplus_{i,j} H^2(U_{ij}) \to H^2(U_{012});\\
C_1 &: \bigoplus_i H^1(U_i) \to \bigoplus_{i,j} H^1(U_{ij}) \to H^1(U_{012});\\
C_0 &: \bigoplus_i H^0(U_i) \to \bigoplus_{i,j} H^0(U_{ij}) \to H^0(U_{012}).
\end{aligned}$$

We can make these complexes a bit more explicit:

$$\begin{array}{ccccccc}
C_3 & : & \mathbb{Z} & \longrightarrow & 0 & \longrightarrow & \mathbb{Z};\\
C_2 & : & \mathbb{Z}_k & \longrightarrow & \mathbb{Z}^3 & \xrightarrow{g_2} & \mathbb{Z}^3;\\
C_1 & : & * & \longrightarrow & \mathbb{Z}^6 & \xrightarrow{g_1} & \mathbb{Z}^3;\\
C_0 & : & \mathbb{Z}^3 & \longrightarrow & \mathbb{Z}^3 & \longrightarrow & \mathbb{Z}.
\end{array}$$

The homology of the complex C_0 is easily computable, and gives

$$H_0(C_0) = H_1(C_0) = H_2(C_0) = 0.$$

In C_1, we claim that the map

$$g_1 : H^1(L_{Y_0},\mathbb{Z}) \oplus H^1(L_{Y_1},\mathbb{Z}) \oplus H^1(L_{X_1},\mathbb{Z}) \to H^1(L_0,\mathbb{Z})$$

is surjective. To see this, take first a basis $\{k_X, k_Y, h\} \subseteq \pi_1(L_0)$, where k_X, k_Y generate, respectively, the kernels of the maps

$$\begin{aligned}
\mathsf{p}_{X*} &: \pi_1(L_0) \to \pi_1(L_{X_1}),\\
\mathsf{p}_{Y*} &: \pi_1(L_0) \to \pi_1(L_{Y_1}),
\end{aligned}$$

and consider the dual basis

$$\{\hat{k}_X,\, \hat{k}_Y, \hat{h}\} \in \hom(\pi_1(L_0),\mathbb{Z}) = H^1(L_0,\mathbb{Z}).$$

We have that $\{\mathsf{p}_{X*}k_Y, \mathsf{p}_{X*}h\} \subseteq \pi_1(L_{X_1})$ and $\{\mathsf{p}_{Y*}k_X, \mathsf{p}_{Y*}h\} \subseteq \pi_1(L_{Y_1})$ are bases as well, and again we can take the dual bases for the comohology

$$\{\widehat{\mathsf{p}_{X*}k_Y}, \widehat{\mathsf{p}_{X*}h}\} \subseteq H^1(L_{X_1}, \mathbb{Z}), \qquad \{\widehat{\mathsf{p}_{Y*}k_X}, \widehat{\mathsf{p}_{Y*}h}\} \subseteq H^1(L_{Y_1}, \mathbb{Z}).$$

It can be easily computed that

$$\begin{aligned}
\hat{k}_X &= g_1(0, -\widehat{\mathsf{p}_{Y*}k_X}, 0),\\
\hat{k}_Y &= g_1(0, 0, \widehat{\mathsf{p}_{X*}k_Y}),\\
\hat{h} &= g_1(0, 0, \widehat{\mathsf{p}_{X*}h}),
\end{aligned}$$

which proves that g_1 is surjective.

Finally we observe that in C_2 the map $\mathbb{Z}_k \to \mathbb{Z}^3$ is necessarily the zero map. With the information we have collected, the second page of the spectral sequence then reads

$$\begin{tikzcd}
\mathbb{Z} \arrow[rrd] & 0 & \mathbb{Z} \\
\mathbb{Z}_k & \ker g_2 & \operatorname{coker} g_2 \\
* & * & 0 \\
\mathbb{Z} & 0 & 0
\end{tikzcd}$$

The arrow in the diagram is the only nonzero differential. Moreover, since $H^4(M, \mathbb{Z}) = H_1(M, \mathbb{Z}) = 0$, the differential is surjective and $\operatorname{coker}(g_2 : \mathbb{Z}^3 \to \mathbb{Z}^3)$ can only be $\mathbb{Z}$, $\mathbb{Z}_m$ or 0, in which cases $\ker g_1 = \mathbb{Z}$, 0 or 0, respectively. In either case, the $E_3 = E_\infty$ page takes one of the following forms:

$$\begin{array}{|ccc}
0 & 0 & \mathbb{Z} \\
\mathbb{Z}_k & \mathbb{Z} & 0 \\
* & * & 0 \\
\mathbb{Z} & 0 & 0 \\
\hline
\end{array}
\qquad\qquad
\begin{array}{|ccc}
\mathbb{Z} & 0 & \mathbb{Z} \\
\mathbb{Z}_k & 0 & 0 \\
* & * & 0 \\
\mathbb{Z} & 0 & 0 \\
\hline
\end{array}$$

In both cases, one can see that $H^3(M, \mathbb{Z}) = H_2(M, \mathbb{Z}) = \mathbb{Z}$. □

8. CURVATURE AND SINGULAR RIEMANNIAN FOLIATIONS BY CIRCLES

In this section we prove Theorem F, which generalizes work of Hsiang and Kleiner [26], in the positively curved case, and of Kleiner [29], and Searle and Yang [46], in the nonnegatively curved case, on simply connected Riemannian 4-manifolds with isometric circle actions.

8.1. **Auxiliary results.**

In this subsection we prove some auxiliary results, without any assumptions on the curvature, and show that there exists a smooth circle action inducing a given singular Riemannian foliation by circles on a compact simply connected smooth 4-manifold. We then prove parts (1) and (2) of Theorem F in Subsection 8.2.

Leaf space structure. We first describe the basic topological properties of the leaf space of a singular Riemannian foliation by circles on a compact, simply connected Riemannian 4-manifold. The leaf space structure is analogous to the orbit space structure of a smooth circle action on a compact, simply connected smooth 4-manifold (cf. [11]).

We begin with the following elemental observation, whose proof follows along the lines of the corresponding statement for smooth actions (cf. [5, Chapter II, Theorem 6.2 and Corollary 6.3]).

Lemma 8.1. *If $(M, \mathcal{F})$ is a singular Riemannian foliation of codimension at least 1, then the fundamental group of M surjects onto the fundamental group of the leaf space M^*.*

We are now ready to describe the structure of the orbit space.

Proposition 8.2. *Let M be a simply connected Riemannian 4-manifold. If M has a singular Riemannian foliation by circles, then the leaf space M^* satisfies the following conditions.*

1. *The leaf space M^* is a simply connected topological 3-manifold, possibly with boundary.*
2. *The components of the 0-dimensional stratum are homeomorphic to 2-spheres or isolated points.*
3. *The boundary components of M^* are 2-sphere components in the 0-dimensional stratum.*

Proof. We first prove (1). By Lemma 8.1, $\pi_1(M)$ surjects onto $\pi_1(M^*)$. Therefore M^* is simply connected. The space of directions Σ_{p^*} at the projection p^* of a singular point $p \in M$ is the quotient of the infinitesimal foliation $\mathcal{F}_p$ of the tangent sphere. This is a 1-dimensional foliation on $\mathbb{S}^3$. If p is an isolated singular point, then the infinitesimal foliation $\mathcal{F}_p$ is regular and, by work of Grove and Gromoll [19], it is a homogeneous foliation, i.e. $\mathcal{F}_p$ is induced by an isometric circle action on $\mathbb{S}^3$. Therefore, the space of directions Σ_{p^*} is homeomorphic to $\mathbb{S}^2$ and p^* is a manifold point in the interior of M^*. If p is a singular point and not isolated, then $\mathcal{F}_p$ is singular and, again, it is homogeneous (see, for example, [43]). Since the foliation is singular, the circle action on $\mathbb{S}^3$ has non-trivial fixed point and it follows that Σ_{p^*} is homeomorphic to a 2-disc. In particular, p^* is a boundary point of M^*. Suppose, finally, that p is a regular point of the foliation. Since $TM_{\mathrm{reg}} = T\mathcal{V} \oplus T\mathcal{H}$ and

TM and $T\mathcal{V}$ are orientable, then $T\mathcal{H}$ is also orientable. Then the holonomy action of $\pi_1(L_p) = \mathbb{Z}$ on the unit normal sphere to L_p, which is isometric to a round unit 2-sphere, preserves orientation. Then $\Sigma_{p^*} = \mathbb{S}^2$ around regular points.

We now prove (2) and (3). Since M^* is a simply connected topological 3-manifold, it is orientable. Using the long exact sequence in homology of the pair $(M^*, \partial M^*)$ and duality, we conclude that $H_1(\partial M^*) = 0$. Since ∂M^* is a disjoint union of surfaces, it follows that the connected components of ∂M^* are 2-spheres. □

Existence of a smooth circle action. We now prove that there exists a smooth circle action inducing a given singular Riemannian foliation by circles.

Theorem 8.3. *A singular Riemannian foliation $(M, \mathcal{F})$ by circles on a compact, smooth simply connected 4-manifold M is induced by a smooth, effective circle action on M.*

Proof. We first prove the following lemma.

Lemma 8.4. *If $(M, \mathcal{F})$ is a 1-dimensional singular Riemannian foliation on a compact, simply connected 4-manifold M, the foliation restricted to the regular part M_{reg} is orientable.*

Proof. Let $F(\mathcal{V})$ be the orthonormal frame bundle of the vertical distribution $\mathcal{V} = T\mathcal{F}$ on M_{reg}. Observe that $F(\mathcal{V}) \to M$ is a $\mathbb{Z}_2$-cover, and we want to prove that $F(\mathcal{V})$ has two connected components. Take an open cover $\{U_0, U_1 \dots U_k\}$ of M_{reg} in the following way. Let Σ_i, $i = 1, \dots, k$ be the components of the 0-dimensional stratum of M. For each Σ_i, let $U_i = B_\epsilon(\Sigma_i) \setminus \Sigma_i$, for some $\epsilon > 0$ small enough so that the sets U_i are disjoint, and let $U_0 = M_{reg} - B_{\epsilon/2}(\cup\Sigma)$. By Proposition 8.2, Σ_i is either a 2-sphere or a point. Therefore, $\mathcal{V}$ is orientable in U_i and $F(\mathcal{V})|_{U_i}$ is disconnected, for $i = 1 \dots k$. Moreover, since U_0 is simply connected, $F(\mathcal{V})|_{U_0}$ is disconnected as well.

Let $A_i = \bigcup_{j=0}^{i} U_i$. We now prove by induction that $F(\mathcal{V})|_{A_i}$ is disconnected or, equivalently, that $\widetilde{H}_0(F(\mathcal{V})|_{A_i}) = \mathbb{Z}_2$, $i = 0, \dots k$. The case $i = 0$ is $\widetilde{H}_0(F(\mathcal{V})|_{U_0}) = \mathbb{Z}_2$, which we have already proved to be true. If $\widetilde{H}_0(F(\mathcal{V})|_{A_i}) = \mathbb{Z}_2$, then by the Mayer-Vietoris sequence applied to the pair $(A_i,\, U_{i+1})$, we have the exact sequence

$$\widetilde{H}_0(F(\mathcal{V})|_{A_i \cap U_{i+1}}) \to \widetilde{H}_0(F(\mathcal{V})|_{A_i}) \oplus \widetilde{H}_0(F(\mathcal{V})|_{U_{i+1}}) \to \widetilde{H}_0(F(\mathcal{V})|_{A_{i+1}}) \to 0.$$

Since the left term is $\mathbb{Z}_2$ and the middle term is $\mathbb{Z}_2 \oplus \mathbb{Z}_2$, the last term must be $\mathbb{Z}_2$. This proves the induction step and, in particular, that

$$\widetilde{H}_0(F(\mathcal{V})_{A_k}) = \widetilde{H}_0(F(\mathcal{V})_{M_{reg}}) = \mathbb{Z}_2,$$

as we wanted. □

By Lemma 8.4, there is a unit vector field X_0' spanning $\mathcal{F}|_{M_{reg}}$. Take now the vector field $X_0(p) = f(p) \cdot X_0'(p)$, where

$$f(p) = |G_p| \cdot \mathrm{length}(L_p)$$

and G_p is the holonomy group (in the foliated sense, cf. [32]) of L_p. Finally, extend X_0 to the singular part, by zeroes. The following proposition shows that the vector field X_0 just defined is smooth, concluding the proof of Theorem 8.3.

Proposition 8.5. *The vector field X_0 defined above is smooth on M.*

Proof. It suffices to prove that f is smooth around the fixed points. Around these points, the infinitesimal foliation is given by a smooth circle action (not necessarily isometric), and

$$f(p) = \int_{\mathbb{S}^1} |g_* X_p^*| dg,$$

where X_p^* is the action field. This is clearly smooth. □

□

By work of Fintushel [11, 12] in combination with work of Pao [37] and Perelman's proof of the Poincaré conjecture [38, 39, 40], a closed, simply connected smooth 4-manifold with a smooth effective circle action is diffeomorphic to a connected sum of copies of $\mathbb{S}^4$, $\pm\mathbb{C}P^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$ (cf. [15] for more details). As a corollary of this result and Theorem 8.3 we obtain the topological classification of compact, simply connected 4-manifolds with a singular Riemannian foliation by circles.

Corollary 8.6. *A compact, simply connected 4-manifold with a singular Riemannian foliation by circles is diffeomorphic to a connected sum of copies of $\mathbb{S}^4$, $\pm\mathbb{C}P^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$.*

8.2. **Proof of (1) and (2) of Theorem F.** By Poincaré duality, $2 \leq \chi(M)$. By Corollary 8.6, to prove parts (1) and (2) of the theorem, it suffices to show that $\chi(M) \leq 3$, when M is positively curved, and $\chi(M) \leq 4$, when M is nonnegatively curved.

By the results in the preceding subsection, the leaf space M^* is a simply connected topological manifold with an Alexandrov space structure. In particular, M^* is positively curved if M has positive curvature, and M^* is nonnegatively curved if M has nonnegative curvature. Since the proof follows as in the proof an isometric circle action, via comparison arguments already found in the literature (cf. [26, 15, 21, 46, 22], we only indicate the necessary steps.

Positively curved case. Suppose first that ∂M^* is not empty, and let F^* be a connected component of ∂M^*. Then, by the Soul Theorem for Alexandrov spaces, there exists a unique point p_0^* at maximal distance from F^* and all the points between F^* and p_0^* must correspond to regular leafs. The point

p_0^* is either a regular leaf or an isolated point in Σ_0. Therefore $\chi(M) \leq 3$. Suppose now that ∂M^* is empty. Then Σ_0 consists only of isolated points. As shown in the proof of Proposition 8.2, the space of directions at an isolated point in the 0-dimensional stratum is isometric to the quotient of a round unit $\mathbb{S}^3$ by an isometric circle action without fixed points. It follows from a triangle comparison argument as in [22] that there can be at most three such points. Therefore, $\chi(\Sigma_0) \leq 3$.

Nonnegatively curved case. Suppose first that ∂M^* has at least two components, ∂M_-^* and ∂M_+^*. Then M^* is isometric to $\partial M_+^* \times [0,1]$ and there are no isolated points in Σ_0. Hence $\chi(M) = 4$. Suppose now that ∂M^* is connected and let C^* be the set at maximal distance from ∂M^* in M^*. There can be at most 2 isolated points in Σ_0 contained in C^*, so $\chi(M^*) \leq 4$. Finally, suppose that ∂M^* is empty and Σ_0 consists only of isolated points. As in the positively curved case, a triangle comparison argument as in [22] implies that there can be at most four such points, so $\chi(M) \leq 4$. □

REFERENCES

1. Alexandrino, M., Rafael B. and Toeben, D., *Progress in the Theory of Singular Riemannian Foliations*, Preprint (2012), arXiv:1203.6829 [math.DG].
2. Alexandrino, M., *Proofs of conjectures about singular Riemannian foliations*, Geom. Dedicata **119** (2006), 219–234.
3. Barden, D., *Simply connected five-manifolds*, Annals of Math. **82** (1965), no. 3, 365–385.
4. Bessieres, L., Besson, G., Boileau, M., Maillot, S., and Porti, J., *Geometrisation of 3-manifolds*, EMS Tracts in Mathematics, volume 13. European Mathematical Society, Zurich, 2010.
5. Bredon, G., *Introduction to compact transformation groups*, Academic Press 48 (1972).
6. Cheeger, J., Fukaya, K. and Gromov, M., *Nilpotent structures and invariant metrics on collapsed manifolds*, J. Amer. Math. Soc. **5** (1992), no. 2, 327–372.
7. Cheeger, J. and Gromov, M., *Collapsing Riemannian manifolds while keeping their curvature bounded. I.*, J. Differential Geom. **23** (1986), no. 3, 309–346.
8. Cheeger, J. and Gromov, M., *Collapsing Riemannian manifolds while keeping their curvature bounded. II.*, J. Differential Geom. **32** (1990), no. 1, 269–298.
9. do Carmo, M., *Riemannian geometry*, Translated from the second Portuguese edition by Francis Flaherty. Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA, 1992.
10. Farrell, F. T., *The Borel conjecture*, Topology of high-dimensional manifolds, No. 1, 2 (Trieste, 2001), 225–298, ICTP Lect. Notes, 9, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2002.
11. Fintushel, R., *Circle actions on simply connected* 4*-manifolds*, Trans. Amer. Math. Soc. **230** (1977), 147–171.
12. Fintushel, R., *Classification of circle actions on* 4*-manifolds*, Trans. Amer. Math. Soc., **242** (1978), 377–390.
13. Freedman, M. H., *The topology of four-dimensional manifolds*, J. Differential Geom. **17** (1982), no. 3, 357–453.
14. Fukaya, K., *Collapsing Riemannian manifolds and its applications*, Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990), 491–500, Math. Soc. Japan, Tokyo, 1991.

15. Galaz-Garcia, F., *Nonnegatively curved fixed point homogeneous manifolds in low dimensions*, Geom. Dedicata **157** (2012), 367–396.
16. Galaz-Garcia, F., *A note on maximal symmetry rank, quasipositive curvature and low-dimensional manifolds*, Preprint (2012), arXiv:1201.1312 [math.DG].
17. Galaz-Garcia, F. and Searle, C., *Low-dimensional manifolds with non-negative curvature and maximal symmetry rank*, Proc. Amer. Math. Soc. **139** (2011), no. 7, 2559–2564.
18. Gorodski, C. and Lytchak, A., *On orbit spaces of representations of compact Lie groups*, J. Reine Angew. Math., to appear. Preprint: arXiv:1109.1739v1 [math.DG].
19. Gromoll, D. and Grove, K., *The low-dimensional metric foliations of Euclidean spheres*, J. Differential Geom. **28** (1988), no. 1, 143–156.
20. Grove, K. and Halperin, S., *Dupin hypersurfaces, group actions and the double mapping cylinder*, J. Differential Geom. **26** (1987), no. 3, 429–459.
21. Grove, K. and Searle, C. *Positively curved manifolds with maximal symmetry-rank*, J. Pure Appl. Algebra **91** (1994), no. 1-3, 137–142.
22. Grove, K. and Wilking, B., *A knot characterization and* 1*-connected nonnegatively curved* 4*-manifolds with circle symmetry*, preprint (2012).
23. Hatcher, A. E., *A proof of the Smale conjecture,* $\mathrm{Diff}(S^3) \simeq \mathrm{O}(4)$, Ann. of Math. (2) **117** (1983), no. 3, 553–607.
24. Hatcher, A. E., *Concordance spaces, higher simple-homotopy theory, and applications*, Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 1, pp. 3–21, Proc. Sympos. Pure Math., XXXII, Amer. Math. Soc., Providence, R.I., 1978.
25. Hsiang, W.C. and Shaneson, J.L., *Fake tori*, 1970 Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969) pp. 18–51 Markham, Chicago, Ill.
26. Hsiang, W.Y. and Kleiner, B., *On the topology of positively curved* 4*-manifolds with symmetry*, J. Differential Geom. **29** (1989), no. 3, 615–621.
27. Kerin, M., *Notes on torus actions on manifolds with quasi-positive curvature*, Preprint.
28. Kim, S. K., McGavran, D. and Pak, J., *Torus group actions on simply connected manifolds*, Pacific J. Math. **53** (1974), 435–444.
29. Kleiner, B., *Riemannian four-manifolds with nonnegative curvature and continuous symmetry*, Ph.D. thesis, University of California, Berkeley, 1990.
30. Kobayashi, S., *Transformation groups in differential geometry*, Reprint of the 1972 edition. Classics in Mathematics. Springer-Verlag, Berlin, 1995.
31. Lytchak, A., *Geometric resolution of singular Riemannian foliations*, Geom. Dedicata **149** (2010), 379–395
32. Molino, P., *Riemannian foliations*, Translated from the French by Grant Cairns. With appendices by Cairns, Y. Carrire, É. Ghys, E. Salem and V. Sergiescu. Progress in Mathematics, 73. Birkhäuser Boston, Inc., Boston, MA, 1988.
33. Oh, H. S., *Toral actions on 5-manifolds*, Trans. Amer. Math. Soc., **278**, no. 1 (1983), 233–252.
34. Orlik, P. and Raymond, F., *Actions of the torus on* 4*-manifolds I*, Trans. Amer. Math. Soc., **152** (1970), 531–559.
35. Orlik, P. and Raymond, F., *Actions of the torus on* 4*-manifolds. II*, Topology **13** (1974), 89–112.
36. Orlik, P. *Seifert manifolds*, Lecture Notes in Mathematics, Vol. 291. Springer-Verlag, Berlin-New York, 1972.
37. Pao, P.S., *Nonlinear circle actions on the* 4*-sphere and twisting spun knots*, Topology **17**, no. 3 (1978), 291–296.
38. Perelman, G., *The entropy formula for the Ricci Flow and its geometric applications*, arXiv:math.DG/0211159, 11 Nov 2002.
39. Perelman, G., *Ricci Flow with surgery on three-manifolds*, arXiv:math.DG/0303109, 10 Mar 2003.

40. Perelman, G., *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv:math.DG/0307245, 17 Jul 2003.
41. Perelman, G. Ya. and Petrunin, A. M., *Extremal subsets in Aleksandrov spaces and the generalized Liberman theorem* (Russian) Algebra i Analiz **5** (1993), no. 1, 242–256; translation in St. Petersburg Math. J. 5 (1994), no. 1, 215–227
42. Püttmann, T. and Searle, C., *The Hopf conjecture for manifolds with low cohomogeneity or high symmetry rank*, Proc. Amer. Math. Soc. **130** (2002), no. 1, 163–166
43. Radeschi, M., *Low dimensional singular Riemannian foliations on spheres*, Ph.D. thesis, University of Pennsylvania (2012).
44. Rong, X., *Positively curved manifolds with almost maximal symmetry rank*, Proceedings of the Conference on Geometric and Combinatorial Group Theory, Part II (Haifa, 2000). Geom. Dedicata **95** (2002), 157–182.
45. Rong, X., *The existence of polarized F-structures on volume collapsed 4-manifolds*, Geom. Funct. Anal. **3** (1993), no. 5, 474–501.
46. Searle, C. and Yang, D., *On the topology of non-negatively curved simply connected 4-manifolds with continuous symmetry*, Duke Math. J. **74** (1994), no. 2, 547–556.
47. Smale, S., *On the structure of* 5*-manifolds*, Ann. of Math. (2) **75** (1962), 38–46.
48. Wilking, B., *Torus actions on manifolds of positive sectional curvature*, Acta Math., **191**, no. 2 (2003), 259–297.
49. Wolak, R. A., *Basic cohomology for singular Riemannian foliations*, Monatsh. Math. **128** (1999), no. 2, 159–163.
50. Wolf, J. A., *Spaces of constant curvature*, Sixth edition. AMS Chelsea Publishing, Providence, RI, 2011.

(Galaz-Garcia) Mathematisches Institut, WWU Münster, Germany.
E-mail address: `f.galaz-garcia@uni-muenster.de`

(Radeschi) Mathematisches Institut, WWU Münster, Germany.
E-mail address: `mrade_02@uni-muenster.de`